

# Stochastic Volterra equations with random functional coefficients in Banach spaces

Alexander Kalinin

Department of Mathematics, LMU Munich

13th Bachelier World Congress  
University of Bologna, July 2026



## The probabilistic framework

---

In what follows, let  $I$  be a non-degenerate interval in  $\mathbb{R}_+$  with  $0 \in I$  and  $E$  be a separable Banach space.

Further, let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space and  $\mathbb{F} = (\mathcal{F}_t)_{t \in I}$  be a filtration of  $\mathcal{F}$  such that the usual conditions hold.

We shall consider the separable Hilbert space  $\ell^2$  of all sequences  $a = (a_i)_{i \in \mathbb{N}}$  of real numbers satisfying  $\sum_{i=1}^{\infty} a_i^2 < \infty$ .

By the [Riesz–Fischer Theorem](#), any infinite-dimensional separable Hilbert space is isometrically isomorphic to  $\ell^2$ .

There is a norm  $|\cdot|_2$  that turns the linear space  $\mathcal{L}_2(\ell^2, E)$  of all  $E$ -valued continuous, linear and **radonifying** operators on  $\ell^2$  into a separable Banach space.

Under the hypothesis that  $E$  is **2-smooth**, for any  $\mathbb{F}$ -progressively measurable process  $U: I \times \Omega \rightarrow \mathcal{L}_2(\ell^2, E)$  satisfying

$$\int_0^\cdot |U_s|_2^2 ds < \infty \quad \text{a.s.}$$

and a sequence  $W = (W^{(i)})_{i \in \mathbb{N}}$  of **independent standard Brownian motions** with respect to  $\mathbb{F}$ , the stochastic integral

$$\int_0^\cdot U_s dW_s$$

can be constructed as an  $E$ -valued continuous  $\mathbb{F}$ -local martingale.

For a sequence  $(Y_i)_{i \in \mathbb{N}}$  of independent  $\mathcal{N}(0, 1)$ -distributed random variables, a continuous linear operator

$$L: \ell^2 \rightarrow E$$

is called **radonifying** if the series  $\sum_{i=1}^{\infty} Y_i L(e_i)$  converges in mean square.

This property is independent of the choice of  $(Y_i)_{i \in \mathbb{N}}$ . Further, if it holds, then the **distribution of  $\sum_{i=1}^{\infty} Y_i L(e_i)$  is uniquely determined.**

It follows that the linear space  $\mathcal{L}_2(\ell^2, E)$ , equipped with the norm

$$|L|_2 := \mathbb{E} \left[ \left| \sum_{i=1}^{\infty} Y_i L(e_i) \right|^2 \right]^{\frac{1}{2}},$$

is a separable Banach space, and  $|L| \leq |L|_2$  for all  $L \in \mathcal{L}_2(\ell^2, E)$ .

### Example (Frobenius norm)

For  $d, m \in \mathbb{N}$  let  $E = \mathbb{R}^m$  and  $A \in \mathbb{R}^{m \times d}$ . If  $L: \ell^2 \rightarrow \mathbb{R}^m$  is given by

$$L(a) := A \begin{pmatrix} a_1 \\ \vdots \\ a_d \end{pmatrix},$$

then  $L \in \mathcal{L}_2(\ell^2, \mathbb{R}^m)$  and

$$\|L\|_2^2 = \mathbb{E}[|AY|^2] = \text{tr}(A^\top A) = |A|^2$$

for the random vector  $Y := (Y_1, \dots, Y_d)^\top$  and the Frobenius norm  $|\cdot|$  on  $\mathbb{R}^{m \times d}$ .

We recall that  $E$  is **2-smooth** if and only if there is a norm  $\|\cdot\|$  equivalent to  $|\cdot|$  such that

$$\|x + y\|^2 + \|x - y\|^2 \leq 2\|x\|^2 + \hat{c}\|y\|^2$$

for all  $x, y \in E$  and some  $\hat{c} \geq 2$ . In this case,  $E$  is **uniformly smooth** and **reflexive**.

If equality holds for  $\hat{c} = 2$ , we recover the **parallelogram identity**, implying that  $\|\cdot\|$  is induced by an inner product.

In general, an equivalent norm to  $|\cdot|$  that turns  $E$  into a Hilbert space does not need to exist, since

$$L^p(S, \mathcal{B}, \mu)$$

is a **2-smooth Banach space** whenever  $(S, \mathcal{B}, \mu)$  is a  $\sigma$ -finite measure space and  $p \geq 2$ .

Within this framework, we consider the **stochastic Volterra equation**

$$X_t = \xi_t + \int_0^t B_{t,s}(X_s) ds + \int_0^t \Sigma_{t,s}(X_s) dW_s \quad \text{a.s.} \quad (1)$$

for  $t \in I$  coupled with a value condition. Here,

- $\xi$  is an  $E$ -valued  $\mathbb{F}$ -progressively measurable process and
- $\mathcal{D}$  is a non-empty set of  $E$ -valued random vectors such that if  $Y \in \mathcal{D}$  and  $Z$  is an  $E$ -valued random vector with

$$Y = Z \quad \text{a.s.}, \quad \text{then} \quad Z \in \mathcal{D}.$$

- Moreover, the **random coefficients**

$$B: I \times I \times \Omega \times \mathcal{D} \rightarrow E \quad \text{and} \quad \Sigma: I \times I \times \Omega \times \mathcal{D} \rightarrow \mathcal{L}_2(\ell^2, E)$$

are **admissible** in a measure theoretical sense.

As Banach-valued functionals,

$$B_{t,s}(\cdot)(\omega) \quad \text{and} \quad \Sigma_{t,s}(\cdot)(\omega)$$

depend on the random vector  $X_s$  of any potential solution  $X$  for any  $s, t \in I$  with  $s \leq t$  and  $\omega \in \Omega$ .

Their domain  $\mathcal{D}$  can be chosen to be linear space

$$\mathcal{L}^p(\Omega, E)$$

of all  $E$ -valued  $p$ -integrable random vectors for  $p \geq 1$ .

Hence, (1) represents a novel class of stochastic Volterra equations for which the presented paper derives unique solutions.

In particular, for controlled and distribution-dependent coefficients we obtain strong solutions.

A **solution** to (1) is an  $E$ -valued  $\mathbb{F}$ -progressively measurable process  $X$  such that

$$X_s \in \mathcal{D} \quad \text{for a.e. } s \in I$$

and the Borel set of all  $t \in I$  for which

$$\int_0^t |B_{t,s}(X_s)| + |\Sigma_{t,s}(X_s)|_2^2 ds < \infty \quad \text{and} \tag{2}$$
$$X_t = \xi_t + \int_0^t B_{t,s}(X_s) ds + \int_0^t \Sigma_{t,s}(X_s) dW_s \quad \text{a.s.}$$

fails has Lebesgue measure zero.

If, moreover,  $X_t \in \mathcal{D}$  and (2) holds for every  $t \in I$ , then  $X$  is called **regular**.

## Controlled distribution-dependent coefficients

Given  $p \geq 1$ , let  $\mathcal{D} = \mathcal{L}^p(\Omega, E)$  and

$$B_{t,s}(X_s) = k(t-s)b(X_s, \alpha_s, \mathcal{L}(X_s, \alpha_s)),$$

$$\Sigma_{t,s}(X_s) = l(t-s)\sigma(X_s, \alpha_s, \mathcal{L}(X_s, \alpha_s))$$

for all  $s, t \in I$  with  $s < t$  and  $X_s \in \mathcal{D}$ , where  $k, l: I \rightarrow \mathbb{R}$  are measurable,

$A$  is a separable Banach space,  $\alpha$  is an  $A$ -valued  $\mathbb{F}$ -progressively measurable control process and the maps

$$b: E \times A \times \mathcal{P}_p(E \times A) \rightarrow E, \quad \sigma: E \times A \times \mathcal{P}_p(E \times A) \rightarrow \mathcal{L}_2(\ell^2, E)$$

are Borel measurable.

For a separable Banach space  $S$ , we recall that  $\mathcal{P}_p(S)$  is the Polish space of all Borel probability measures on  $S$  satisfying

$$\int_S |x|^p \mu(dx) < \infty,$$

equipped with the  $p$ th Wasserstein metric given by

$$\vartheta_p(\mu, \nu) := \inf_{\theta \in \mathcal{P}(\mu, \nu)} \left( \int_{S \times S} |x - y|^p d\theta(x, y) \right)^{\frac{1}{p}},$$

where  $\mathcal{P}(\mu, \nu)$  is the convex space of all Borel probability measures  $\theta$  on  $S \times S$  that admit

$$\mu \quad \text{and} \quad \nu$$

as first and second marginal distributions, respectively.

Moreover, **strong solutions** to (1) can be defined using the natural filtration of  $\xi$ ,  $\alpha$  and  $W^{(i)}$ , where  $i \in \mathbb{N}$ .

## Unique strong solutions (K., 2026)

Let  $p \geq 2$ ,  $k$  and  $l^2$  be locally integrable and  $b$  and  $\sigma$  be Lipschitz continuous. Then the following assertions hold:

- (i) If  $\text{ess sup}_{s \in [0, t]} \mathbb{E}[|\xi_s|^p + |\alpha_s|^p] < \infty$  for all  $t \in I$ , then there exists a **unique strong solution**  $X^\xi$  to (1) such that

$$\text{ess sup}_{s \in [0, t]} \mathbb{E}[|X_s^\xi|^p] < \infty \quad \text{for all } t \in I.$$

- (ii) If merely  $\int_0^t \mathbb{E}[|\xi_s|^p + |\alpha_s|^p] ds < \infty$  for any  $t \in I$ , then we obtain a **unique strong solution**  $X^\xi$  satisfying

$$\int_0^t \mathbb{E}[|X_s^\xi|^p] ds < \infty \quad \text{for any } t \in I.$$

## Unique strong solutions (cont.)

(iii) Let  $\xi = \xi_0$  a.s. and  $\mathbb{E}[|\xi_0|^q] + \text{ess sup}_{s \in [0, t]} \mathbb{E}[|\alpha_s|^q] < \infty$  for all  $q \geq 2$  and  $t \in I$ . Further, assume that

$$k(u) = u^{\gamma-1} \quad \text{and} \quad l(u) = u^{\delta-\frac{1}{2}}$$

for all  $u \in I \setminus \{0\}$ , where  $\gamma \in ]0, 1]$  and  $\delta \in ]0, \frac{1}{2}]$ . Then (1) admits a **unique, strong and regular solution**  $X^{\xi_0}$  such that

$$\sup_{s \in [0, t]} \mathbb{E}[|X_s^{\xi_0}|^q] < \infty$$

for all  $t \in I$  and  $q \geq 2$  and whose paths are locally  $\beta$ -Hölder continuous for any  $\beta \in ]0, \gamma \wedge \delta[$ .

To establish the existence of solutions, **without requiring a priori any path regularity**, we use the progressive  $\sigma$ -field  $\mathcal{A}$  on  $I \times \Omega$  as follows.

## Progressively measurable weak modifications of stochastic Volterra integrals (K., 2026)

Let  $U: I \times I \times \Omega \rightarrow \mathcal{L}_2(\ell^2, E)$  be a  $\mathcal{B}(I) \otimes \mathcal{A}$ -measurable map such that

$$\int_0^t |U_{t,s}|_2^2 ds < \infty \quad \text{a.s. for any } t \in I.$$

Then there exists an  $E$ -valued  $\mathbb{F}$ -progressively measurable process  $X$  such that

$$X_t = \int_0^t U_{t,s} dW_s \quad \text{a.s. for a.e. } t \in I.$$

The fact that solutions can be chosen to be **strong** relies on a general measurability result.

## Measurability in the $p$ th Wasserstein space (K., 2026)

Let  $I$  be merely a non-empty set, endowed with a  $\sigma$ -field  $\mathcal{I}$ , and  $S$  be a separable Banach space. Then for any product measurable process  $X: I \times \Omega \rightarrow S$  such that

$$\mathbb{E}[|X_t|^p] < \infty$$

for all  $t \in I$ , the distribution map

$$I \rightarrow \mathcal{P}_p(S), \quad t \mapsto \mathcal{L}(X_t)$$

is Borel measurable.

# References

---

- [1] A. Kalinin.  
Stochastic Volterra equations with random functional coefficients in Banach spaces.  
*arXiv preprint arXiv:2602.09922*, 2026.
- [2] D. J. Prömel and D. Scheffels.  
Mean-field stochastic Volterra equations.  
*Journal of Applied Probability*, pages 1–25, 2026.
- [3] A. Kalinin.  
Resolvent and Gronwall inequalities and fixed points of evolution operators.  
*arXiv preprint arXiv:2412.20764*, 2024.
- [4] J. M. A. M. van Neerven, M. C. Veraar, and L. Weis.  
Stochastic integration in UMD Banach spaces.  
*Ann. Probab.*, 35(4):1438–1478, 2007.

**Thank you for your attention!**